



Process Unit Overview

The tables below summarize the current composition of the process unit and associated drift cell. The process unit is pictured to the left.

Process Unit Size

Drainage Area	14 km ²
Nearshore Zone	6.6 km ²
Shoreline Length	8.4 km

Drift Cell Composition

Divergence	3%
Transport	64%
Convergence	0%
No Appreciable Drift	33%

Nearshore Process Degradation Summary

Degradation categories reflect the extent to which anthropogenic stressors affect 11 nearshore processes, in both current and future scenarios, relative to other process units. An overall degradation group assignment is made (none, least, less, moderate, more and most) for each scenario. High degradation may indicate areas suitable for restoration while low degradation areas may be more suited towards protection. Future degradation conditions were projected into 2060 assuming current land development trends. See publication 2011-02 (Schlenger et al) for more information.

Nearshore Process	Degradation	
	Current	Future
Sediment Input	Medium	Medium
Sediment Transport	Low	Low
Sediment Accretion	Low	Low
Tidal Flow	Low	Low
Distributary Channels*	N/A	N/A
Tidal Channels	None	Low
Freshwater Input	Medium	Medium
Detritus Import and Export	Medium	Medium
Exchange of Aquatic Organisms	Low	Low
Physical Disturbance*	Medium	Medium
Solar Radiation	Low	Low
Overall Degradation	Less	Less

*Distributary Channels Process does not apply in Shoreline Process Units.
Physical Disturbance Process does not apply in Delta Process Units.

Recommended Management Strategy

Table displays a preliminary management recommendation (protect, restore, or enhance) for landforms within the process unit, based on current degradation and potential to provide ecosystem goods and services. See publication 2011-04 (Cereghino et al) for more information.

Landform	Degradation	Potential	Recommendation
Barrier Embayments	Medium	High	Protect High
Beaches	Medium	Medium	Restore Low

Shoreline Alterations Historic (ca 1850-1880) and Current (ca 2000-2009)

Tables to the right show itemized changes to historic attributes such as tidal wetland areas and anthropogenic modifications (stressors) along the shore. See publication 2011-01 (Simenstad et al) for more information.

Upland Alterations

Modifications found upland of shoreline. Area within 200 feet of shoreline is considered adjacent upland. Watershed area includes adjacent upland.

Modification	Adjacent Upland	Watershed Area
Fish Passage Barrier	1	15
Stream Crossings	2	8
Dams	0	0

Tidal Wetland	Historic Area (sq km)	Current Area (sq km)	Change
Tidal Freshwater	0	0	No Change
Oligohaline Transition	0	0	No Change
Estuarine Mixing	0.14	0	Loss (-100% Area)
Euryhaline Unvegetated	0	0	No Change
Totals	0.14	0	

Shoreline Stressor	Extent of Stressor in Process Unit
Abandoned Railroads	0 km (0% of Shoreline Length)
Active Railroads	0 km (0% of Shoreline Length)
Armoring	1.8 km (22% of Shoreline Length)
Roads	0 km (< 1% of Shoreline Length)
Tidal Barriers	0 km (0% of Shoreline Length)
Breakwater/Jetties	0 Count comprising 0 km of Shoreline
Marinas	0 Count comprising 0 sq km
Nearshore Fill	0.01 sq km
Overwater Structures	7 Count comprising 0.002 sq km

Landform Composition

Historic (ca 1850-1880) and Current (ca 2000-2009)

Landforms are a shoreline classification system based on geomorphic processes (see publication 2006-02 [Shipman]). Differences between historic and current conditions are assumed to be due to shoreline alterations that result in landform change, but could also represent error in mapping capabilities. See publication 2011-01 (Simenstad et al) for more information.

Landforms in Process Unit	Landform Count			Landform Length (km)		
	Historic	Current	Change	Historic	Current	Change (%)
Bluff-Backed Beach	3	3	None	4.8	4.6	-4
Barrier Beach	3	2	-1	0.3	0.3	-6
Barrier Estuary	2	1	-1	4.3	3.0	-31
Barrier Lagoon	1	0	-1	0.2	0.0	-100
Artificial	0	1	1	0.0	0.5	100